

Development and applications of GeoChip 3.0 for analysis of microbial community structures, compositions, and potential functions

Z. He^{1,2}, Y. Deng^{1,2}, J. D. Van Nostrand^{1,2}, **L. Wu**^{1,2}, C. L. Hemme^{1,2}, T. J. Gentry³, J. Liebich⁴, Q. Tu¹, A. P. Arkin^{5,2}, T. C. Hazen^{5,2}, J. Zhou^{1,2};

¹The Univ. of Oklahoma, Norman, OK, ²Virtual Inst. for Microbial Stress and Survival, [Http://vimss.lbl.gov](http://vimss.lbl.gov), CA, ³Texas A&M Univ., College Station, TX, ⁴Forschungszentrum Jülich GmbH, Jülich, GERMANY, ⁵Lawrence Berkeley Natl. Lab., Berkeley, CA

Acknowledgment

This work was part of the Virtual Institute for Microbial Stress and Survival (<http://VIMSS.lbl.gov>) supported by the U. S. Department of Energy, Office of Science, Office of Biological and Environmental Research, Genomics Program:GTL through contractDE-AC02-05CH11231 between Lawrence Berkeley National Laboratory and the U. S. Department of Energy.

Abstract: Functional gene arrays (FGAs), or GeoChip constructed with key genes involved in various biological and geochemical processes have been widely used to analyze microbial communities. Based on GeoChip 2.0, a new generation of GeoChip (GeoChip 3.0) has been developed. GeoChip 3.0 has several new features in terms of coverage, design, and data analysis. GeoChip 3.0 covers approximately 50,000 gene sequences for 306 gene families, and especially functional genes involved in antibiotic resistance and energy metabolism, and phylogenetic marker GyrB have been added, which allows us to obtain more information about microbial communities and analyze more diverse environmental samples. For probe design, sequence-specific, exclusive group-specific, inclusive group-specific, and quasi-group probes with different specificities were selected, and all those probes together greatly increase the gene coverage. For data analysis, three strategies have been implemented. First, a universal standard has been implemented so that data normalization and comparison of samples from different sites, time points, or laboratories can be conducted. Second, a genomic standard is used to quantitatively analyze gene abundance. Third, a software package (including databases) has been developed for sequence retrieval, probe design, information storage, and especially, data analysis and automatic update, which greatly facilitate the management and analysis of complicated GeoChip data sets. GeoChip 3.0 has been used for analyses of responses of microbial communities to elevated CO₂ and increased temperature. The results showed that autotrophic CO₂ and N₂ fixation by microorganisms increased significantly in response to elevated CO₂. This also demonstrates that GeoChip can provide insights into biogeochemical processes and functional activities of microbial communities important to human health, agriculture, energy, global climate change, ecosystem management, and environmental cleanup and restoration. It is also particularly useful for providing direct linkages of microbial genes/populations to ecosystem processes and functions.

Keyword (Complete): microarray ; functional gene ; microbial community